

THE COHOMOLOGY OF THE MOTIVIC STEENROD ALGEBRA OVER $\mathrm{Spec} \mathbb{C}$

DANIEL C. ISAKSEN

ABSTRACT. The purpose of this article is to compute the cohomology of the motivic Steenrod algebra over $\mathrm{Spec} \mathbb{C}$ through the geometric 70-stem. The main computational tool is the motivic May spectral sequence. Everywhere in this article, we are working only over $\mathrm{Spec} \mathbb{C}$, and we are always computing at the prime $p = 2$.

1. INTRODUCTION

This article is one piece of a long-term project to explore stable homotopy groups in motivic homotopy theory. Just as in classical homotopy theory, the stable motivic homotopy groups are key to understanding the behavior of the stable cellular motivic homotopy category. The goal of this long-term project is to supply a large quantity of computational data about motivic stable homotopy rings. This data can be used to detect large-scale patterns in motivic stable homotopy that do not arise in classical stable homotopy. Even better, the richer motivic stable homotopy setting allows one to easily detect some otherwise difficult facts about the classical situation. We provide some examples of these payoffs in Section 1.2.

One important difference between the classical case and the motivic case is that not every motivic spectrum is built out of spheres, i.e., not every motivic spectrum is cellular. Stable cellular motivic homotopy theory is more tractable than the full motivic homotopy theory, and many motivic spectra of particular interest, such as the Eilenberg-Mac Lane spectrum $H\mathbb{F}_2$, the algebraic K -theory spectrum KGL , and the algebraic cobordism spectrum MGL , are all cellular. On the other hand, some arithmetic and algebro-geometric questions are not accessible via the cellular theory.

Although one can study motivic homotopy theory over any base field (or even more general base schemes), we will work only over $\mathrm{Spec} \mathbb{C}$, or any algebraically closed field of characteristic 0. The point is to explore the parts of motivic homotopy theory that don't depend on the subtle arithmetic of the base field. Even in this simplest case, we can find exotic phenomena that have no classical analogues.

One way to approach the stable motivic homotopy groups is through the motivic Adams spectral sequence. First we must choose a prime p . In this article, we are always working at $p = 2$. In order to carry out the 2-primary motivic Adams spectral sequence, one must first obtain the E_2 -term, which is the cohomology of the motivic Steenrod algebra. The purpose of this article is to describe this E_2 -term explicitly through the geometric 70-stem. We will use the motivic May spectral sequence to carry out the computation.

After obtaining the E_2 -term, the next step is to analyze differentials in the motivic Adams spectral sequence. This will appear in a sequel [12] to this article.

A second purpose of this calculation is to provide input to the ρ -Bockstein spectral sequence for computing the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{R}$. The analysis of the ρ -Bockstein spectral sequence, and the further analysis of the motivic Adams spectral sequence over $\text{Spec } \mathbb{R}$, will appear in [DHI].

The exposition of such a technical calculation creates some inherent challenges. The goal of the article is not to describe every step from first principles. Rather, it is a guide for reproducing the computation for those who are already well-versed in computations with the classical May spectral sequence [Ma1] [T]. We give the key facts and examples of how the arguments work.

This article is a natural sequel to [DI], where the first computational properties of the motivic May spectral sequence, as well as of the motivic Adams spectral sequence, were established. We will review the key inputs from [DI].

We have been careful to adhere to a philosophy of logical dependence. We will freely borrow results from the classical May spectral sequence, i.e., from [Ma1] and [T], without further explanation. The point is that this article primarily focuses on the phenomena that are truly new in the motivic context.

We will also need some facts from the cohomology of the classical Steenrod algebra that have been verified only by machine [Br1] [Br2]. The machine computations construct a minimal resolution of $\mathbb{F}_2$ over the classical Steenrod algebra. From this resolution, one can derive the full structure of the cohomology of the classical Steenrod algebra, including multiplicative structures, Massey products, and squaring operations.

1.1. Outline. We begin in Section 2 with a review of the basic facts about the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$, as well as the motivic May spectral sequence over $\text{Spec } \mathbb{C}$. We also establish the computational relationship between the motivic setting and the classical setting. This section is mostly a summary of some parts of [DI].

Next, in Section 3 we describe the main points in computing the motivic May spectral sequence through the geometric 70-stem. We rely heavily on results of [Ma1] and [T], but we must also compute several exotic differentials, i.e., differentials that do not occur in the classical situation.

Having obtained the E_∞ -term of the motivic May spectral sequence, the next step is to consider hidden extensions. In Section 4, we are able to resolve every possible hidden extension by τ , h_0 , h_1 , and h_2 through the range that we are considering, i.e., up to the geometric 70-stem. The primary tools here are:

- (1) shuffling relations among Massey products.
- (2) squaring operations on Ext groups in the sense of [Ma2].
- (3) classical hidden extensions established by machine computation [Br1].

In Section 5, we consider what happens when the element h_1 is inverted in the cohomology of the motivic Steenrod algebra. Because h_1^k is non-zero for all k , this becomes an interesting calculation.

Section 6 consists of a series of tables that are essential for bookkeeping throughout the computations. The most important of these is Table 14, which lists the multiplicative generators of the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$.

Finally, in Section 7, we provide charts that describe the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$, through the geometric 70-stem. These charts

are the main contribution of this article. The diagrams are best viewed in color, although they are legible in black and white. Large format contiguous charts are available as well.

1.2. Some examples. In this section, we describe several of the computational intricacies that are established later in the article. We also present a few questions deserving of further study. On a first reading, this section may be safely skipped and reviewed later.

Example 1.3. An obvious question, which already arose in [DI], is to find elements that are killed by τ^n but not by τ^{n-1} , for various values of n .

The element h_2g^2 , which is multiplicatively indecomposable, is the first example of an element that is killed by τ^3 but not by τ^2 . This occurs because of a hidden extension $\tau \cdot \tau h_2g^2 = Ph_1^4h_5$. There is an analogous relation $\tau^2h_2g = Ph_4$, which occurs already in the May spectral sequence. We do not know if this generalizes to a family of relations of the form $\tau^2h_2g^{2^k} = Ph_1^{2^{k+2}-4}h_{k+4}$.

We will show in [I2] that h_2g^2 represents an element in motivic stable homotopy that is killed by τ^3 but not by τ^2 . This requires an analysis of the motivic Adams spectral sequence. In the vicinity of g^{2^k} , one might hope to find elements that are killed by τ^n but not by τ^{n-1} , for large values of n .

Example 1.4. Classically, there is a relation $h_3 \cdot e_0 = h_1h_4c_0$ in the geometric 24-stem of the cohomology of the Steenrod algebra. This relation is hidden on the E_∞ -page of the May spectral sequence. We now give a proof of this classical relation that uses the cohomology of the motivic Steenrod algebra. It still relies on a hidden extension, but in a more elementary way.

Motivically, it turns out that $h_2^3e_0$ is non-zero, even though it is zero classically. This follows from the hidden extension $h_0 \cdot h_2^2g = h_1^3h_4c_0$ (see Lemma 4.11). The relation $h_2^3 = h_1^2h_3$ then implies that $h_1^2h_3e_0$ is non-zero. Therefore, h_3e_0 is non-zero as well, and the only possibility is that $h_3e_0 = h_1h_4c_0$.

Example 1.5. Notice the hidden extension $h_0 \cdot h_2^2g^2 = h_1^7h_5c_0$ (and similarly, the hidden extension $h_0 \cdot h_2^2g = h_1^3h_4c_0$ that we discussed above in Example 1.4).

The next example in this family is $h_0 \cdot h_2^2g^3 = h_1^9D_4$, which at first does not appear to fit a pattern. However, there is a hidden extension $c_0 \cdot i_1 = h_1^4D_4$, so we have $h_0 \cdot h_2^2g^3 = h_1^5c_0i_1$. Presumably, there is an infinitely family of hidden extensions in which $h_0 \cdot h_2^2g^k$ equals some power of h_1 times c_0 times an element related to Sq^0 of elements associated to the image of J .

It is curious that $c_0 \cdot i_1$ is divisible by h_1^4 . An obvious question for further study is to determine the h_1 -divisibility of c_0 times elements related to Sq^0 of elements associated to the image of J . For example, what is the largest power of h_1 that divides g^2i_1 ?

Example 1.6. Beware that g^2 and g^3 are not actually elements of the geometric 40-stem and 60-stem respectively. Rather, it is only τg^2 and τg^3 that exist (similarly, g does not exist in the 20-stem, but τg does exist). The reason is that there are May differentials taking g^2 to $h_1^8h_5$, and g^3 to $h_1^6i_1$. In other words, τg^2 and τg^3 are multiplicatively indecomposable elements. More generally, we anticipate that the element g^k does not exist because it supports a May differential related to Sq^0 of an element in the image of J .

Example 1.7. There is an isomorphism from the cohomology of the classical Steenrod algebra to the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$ concentrated in degrees of the form $(2s + f, f, \frac{s+f}{2})$. This isomorphism preserves all higher structure, including squaring operations and Massey products. See Section 2.9 for more details.

For example, the existence of the classical element Ph_2 immediately implies that h_3g must be non-zero in the motivic setting; no calculations are necessary.

Another example is that $h_1^{2^k-1}h_{k+2}$ is non-zero motivically for all $k \geq 1$, because $h_0^{2^k-1}h_{k+1}$ is non-zero classically.

Example 1.8. Many elements are h_1 -local in the sense that they support infinitely many multiplications by h_1 . In fact, any product of the symbols h_1 , c_0 , P , d_0 , e_0 , and g , if it exists, is non-zero. This is detectable in the cohomology of motivic $A(2)$ [I1].

Moreover, the element B_1 in the 46-stem is h_1 -local, and any product of B_1 with elements in the previous paragraph is again h_1 -local. This is detectable in the h_1 -local cohomology of motivic $A(3)$, as described in Theorem 5.6. We will see in [I2] that these h_1 -local computations lead to interesting results about classical Adams differentials.

The h_1 -localization of the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$ deserves further study. It should be possible to make a full calculation in a much larger range than is possible for the unlocalized cohomology of the motivic Steenrod algebra. Since localization is exact, one may localize h_1 in the E_1 -term of the May spectral sequence. This makes the E_1 -term much smaller, although still complicated in sufficiently high degrees.

Example 1.9. The motivic analogue of “wedge” subalgebra [MT] appears to be more complicated than the classical version. For example, none of the wedge elements support multiplications by h_0 in the classical case. Motivically, many wedge elements do support h_0 multiplications. The results in this paper naturally call for further study of the structure of the motivic wedge.

1.10. Acknowledgements. The author thanks R. R. Bruner for the use of his extensive library of machine-assisted classical computations. Many of the results in this article would have been impossible to discover without the guidance of this data. Similarly, the author thanks D. Dugger, whose machine-assisted motivic computations give the $\mathbb{M}_2$ -module structure of the cohomology of the motivic Steenrod algebra.

M. Tangora provided invaluable help with various technical computations in the classical May spectral sequence.

M. Mahowald and M. Behrens played especially pivotal roles in giving the author the courage to attempt the calculation.

2. FOUNDATIONS

Stable motivic homotopy is bigraded. In a typical bigraded (p, q) , we refer to p as the geometric stem, to $p - q$ as the topological stem, and to q as the weight.

We will use the following notation extensively:

- (1) $\mathbb{M}_2$ is the mod 2 motivic cohomology of $\text{Spec } \mathbb{C}$.
- (2) A is the mod 2 motivic Steenrod algebra over $\text{Spec } \mathbb{C}$.

- (3) Ext is the trigraded ring $\text{Ext}_A(\mathbb{M}_2, \mathbb{M}_2)$.
- (4) A_{cl} is the classical mod 2 Steenrod algebra.
- (5) Ext_{cl} is the bigraded ring $\text{Ext}_{A_{\text{cl}}}(\mathbb{F}_2, \mathbb{F}_2)$.

The following two deep theorems of Voevodsky are the starting points of our calculations.

Theorem 2.1 ([V1]). $\mathbb{M}_2$ is the bigraded ring $\mathbb{F}_2[\tau]$, where τ has bidegree $(0, 1)$.

Theorem 2.2 ([V2] [V3]). The motivic Steenrod algebra A is the $\mathbb{M}_2$ -algebra generated by elements Sq^{2k} and Sq^{2k-1} for all $k \geq 1$, of bidegrees $(2k, k)$ and $(2k-1, k-1)$ respectively, and satisfying the following relations for $a < 2b$:

$$\text{Sq}^a \text{Sq}^b = \sum_c \binom{b-1-c}{a-2c} \tau^? \text{Sq}^{a+b-c} \text{Sq}^c.$$

The symbol $?$ stands for either 0 or 1, depending on which value makes the formula balanced in weight. See [DI] for a more detailed discussion of the motivic Adem relations.

The A -module structure on $\mathbb{M}_2$ is trivial, i.e., every Sq^k acts by zero. This follows for simple degree reasons.

2.3. Ext groups. We are interested in computing $\text{Ext}_A(\mathbb{M}_2, \mathbb{M}_2)$, which we abbreviate as Ext . This is a trigraded object. We will consistently use degrees of the form (s, f, w) , where:

- (1) f is the Adams filtration, i.e., the homological degree.
- (2) $s+f$ is the internal geometric degree, i.e., corresponds to the first coordinate in the bidegrees of A .
- (3) s is the geometric stem, i.e., the internal geometric degree minus the Adams filtration.
- (4) w is the weight.

Note that $\text{Ext}^{*,0,*} = \text{Hom}_A^{*,*}(\mathbb{M}_2, \mathbb{M}_2)$ is dual to $\mathbb{M}_2$. We will abuse notation and write $\mathbb{M}_2$ for this dual. Beware that now τ , which is really the dual of the τ that we discussed earlier, has degree $(0, 0, -1)$. Since Ext is a module over $\text{Ext}^{*,0,*}$, i.e., over $\mathbb{M}_2$, we will always describe Ext as an $\mathbb{M}_2$ -module.

The following result is the key tool for comparing classical and motivic computations. The point is that the motivic and classical computations become the same after inverting τ .

Proposition 2.4 ([DI]). *There is an isomorphism of rings*

$$\text{Ext} \otimes_{\mathbb{M}_2} \mathbb{M}_2[\tau^{-1}] \cong \text{Ext}_{A_{\text{cl}}} \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau, \tau^{-1}].$$

2.5. The motivic May spectral sequence. The classical May spectral sequence arises by filtering the classical Steenrod algebra by powers of the augmentation ideal. The same approach can be applied in the motivic setting to obtain the motivic May spectral sequence. Details appear in [DI]. Next we review the main points.

The motivic May spectral sequence is quadruply graded. We will always use gradings of the form (m, s, f, w) , where m is the May filtration, and the other coordinates are as explained earlier.

Let $\text{Gr}(A)$ be the associated graded algebra of A , and let $\text{Gr}(A_{\text{cl}})$ be the associated graded algebra of A_{cl} .

Theorem 2.6. *The motivic May spectral sequence takes the form*

$$E_2 = \text{Ext}_{\text{Gr}(A)}^{(m,s,f,w)}(\mathbb{M}_2, \mathbb{M}_2) \Rightarrow \text{Ext}_A^{(s,f,w)}(\mathbb{M}_2, \mathbb{M}_2).$$

Remark 2.7. As in the classical May spectral sequence, the odd differentials must be trivial for degree reasons.

Proposition 2.8. *After inverting τ , there is an isomorphism of spectral sequences between the motivic May spectral sequence of Theorem 2.6 and the classical May spectral sequence, tensored over $\mathbb{F}_2$ with $\mathbb{F}_2[\tau, \tau^{-1}]$.*

Proof. Start with the fact that $A[\tau^{-1}]$ is isomorphic to $A_{\text{cl}} \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau, \tau^{-1}]$, with the same May filtrations. $\square$

This proposition means that differentials in the motivic May spectral sequence must be compatible with the classical differentials. This fact is critical to the success of our computations.

2.9. Ext in Chow degree zero.

Definition 2.10. *Let A' be the subquotient $\mathbb{M}_2$ -algebra of A generated by Sq^{2k} for all $k \geq 0$, subject to the relation $\tau = 0$.*

Lemma 2.11. *There is an isomorphism $A_{\text{cl}} \rightarrow A'$ that takes Sq^k to Sq^{2k} .*

The isomorphism takes elements of degree n to elements of bidegree $(2n, n)$.

Proof. Modulo τ , the motivic Adem relation for $\text{Sq}^{2a} \text{Sq}^{2b}$ takes the form

$$\text{Sq}^{2a} \text{Sq}^{2b} = \sum_c \binom{2b-1-2c}{2a-4c} \text{Sq}^{2a+2b-2c} \text{Sq}^2 c.$$

A standard fact from combinatorics says that

$$\binom{2b-1-2c}{2a-4c} = \binom{b-1-c}{a-2c}$$

modulo 2. $\square$

Definition 2.12. *Let M be a bigraded A -module. The Chow degree of an element m in degree (t, w) is equal to $t - 2w$.*

The terminology arises from the fact that the Chow degree is fundamental in Bloch's higher Chow group perspective on motivic cohomology [Bl].

Definition 2.13. *Let M be an A -module. Define the A' -module $\text{Ch}_0(M)$ to be the subset of M consisting of elements of Chow degree zero, with A' -module structure induced from the A -module structure on M .*

The A' -module structure on $\text{Ch}_0(M)$ is well-defined since Sq^{2k} preserves Chow degrees.

Theorem 2.14. *There is an isomorphism from Ext_{cl} to the subalgebra of Ext consisting of elements whose internal Chow degree is zero. This isomorphism takes classical elements of degree (s, f) to motivic elements of degree $(2s + f, f, \frac{s+f}{2})$, and it preserves all higher structure, including products, squaring operations, and Massey products.*

Remark 2.15. The internal Chow degree of an element of $\text{Ext}^{s,f,w}$ is $s + f - 2w$, since the internal geometric degree of such an element is $s + f$.

Proof. There is a natural transformation $\text{Hom}_A(-, \mathbb{M}_2) \rightarrow \text{Hom}_{A'}(C_0(-), \mathbb{F}_2)$, since $\text{Ch}_0(\mathbb{M}_2) = \mathbb{F}_2$. Since Ch_0 is an exact functor, the derived functor of the right side is $\text{Ext}_{A'}(\text{Ch}_0(-), \mathbb{F}_2)$. The universal property of derived functors gives a natural transformation $\text{Ext}_A(-, \mathbb{M}_2) \rightarrow \text{Ext}_{A'}(\text{Ch}_0(-), \mathbb{F}_2)$. Apply this natural transformation to $\mathbb{M}_2$ to obtain $\text{Ext}_A(\mathbb{M}_2, \mathbb{M}_2) \rightarrow \text{Ext}_{A'}(\text{Ch}_0(\mathbb{M}_2), \mathbb{F}_2)$. The left side is Ext , and the right side is isomorphic to Ext_{cl} since A' is isomorphic to A_{cl} .

We have now obtained a map $\text{Ext} \rightarrow \text{Ext}_{\text{cl}}$. To verify that this map is an isomorphism on the Chow degree zero part of Ext , compare the classical May spectral sequence with the part of the motivic May spectral sequence in Chow degree zero. The motivic elements h_{i0} have Chow degree 1, while the motivic elements h_{ij} have Chow degree 0 for $j > 0$. It follows that the motivic E_1 -term in Chow degree zero is the polynomial algebra over $\mathbb{F}_2$ generated by h_{ij} for $i > 0$ and $j > 0$. This is isomorphic to the classical E_1 -term, where the motivic element h_{ij} corresponds to the classical element $h_{i,j-1}$. $\square$

Remark 2.16. Similar methods show that Ext is concentrated in positive Chow degree. The map $\text{Ext} \rightarrow \text{Ext}_{\text{cl}}$ constructed in the proof takes elements in strictly positive Chow degree to zero. Thus, the quotient of Ext by the strictly positive Chow degree part is isomorphic to Ext_{cl} .

3. COMPUTING WITH THE MOTIVIC MAY SPECTRAL SEQUENCE

3.1. The E_1 -term. The E_2 -term of the May spectral sequence is the cohomology of an algebra. In other words, the May spectral sequence really starts with an E_1 -term. As described in [DI], the motivic E_1 -term is essentially the same as the classical E_1 -term. Specifically, the motivic E_1 -term is a polynomial algebra over $\mathbb{M}_2$ with generators h_{ij} for all $i > 0$ and $j \geq 0$, where:

- (1) h_{i0} has degree $(i, 2^i - 2, 1, 2^{i-1} - 1)$.
- (2) h_{ij} has degree $(i, 2^j(2^i - 1) - 1, 1, 2^{j-1}(2^i - 1))$ for $j > 0$.

The d_1 -differential is described by the formula:

$$d_1(h_{ij}) = \sum_{0 < k < i} h_{kj} h_{i-k, k+j}.$$

3.2. The E_2 -term. We now describe the E_2 -term of the motivic May spectral sequence. As explained in [DI], it turns out that the motivic E_2 -term is essentially the same as the classical E_2 -term. The following proposition makes this precise.

Proposition 3.3 ([DI]). *There are graded ring isomorphisms*

- (a) $\text{Gr}(A) \cong \text{Gr}(A_{\text{cl}}) \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau]$.
- (b) $\text{Ext}_{\text{Gr}(A)}(\mathbb{M}_2, \mathbb{M}_2) \cong \text{Ext}_{\text{Gr}(A_{\text{cl}})}(\mathbb{F}_2, \mathbb{F}_2) \otimes_{\mathbb{F}_2} \mathbb{F}_2[\tau]$.

In other words, explicit generators and relations for the E_2 -term can be lifted directly from the classical situation [T].

Moreover, because of Proposition 2.8, the values of d_2 can also be lifted from the classical situation, except that a few factors of τ show up to give the necessary weights. For example, classically we have the differential

$$d_2(b_{20}) = h_1^3 + h_0^2 h_2.$$

Motivically, this does not make sense, since b_{20} and $h_0^2 h_2$ have weight 2, while h_1^3 has weight 3. Therefore, the motivic differential must be

$$d_2(b_{20}) = \tau h_1^3 + h_0^2 h_2.$$

Table 1 lists the multiplicative generators of the E_2 -term through the geometric 70-stem, and Table 2 lists a generating set of relations for the E_2 -term in the same range. Table 1 also gives the values of the May d_2 differential, all of which are easily deduced from the classical situation [T].

3.4. The E_4 -term. Although the E_2 -term is quite large, the d_2 differential is also very destructive. As a result, the E_4 -page becomes manageable. We obtain the E_4 -page by direct computation with the d_2 differential.

The multiplicative generators for the E_4 -term through the geometric 70-stem break into two groups. The first group consists of generators that survive to E_∞ and become multiplicative generators of Ext; these are listed in Table 14 as elements that occur on or before the E_4 -page. The second group consists of generators that do not survive to E_∞ ; these are listed in Table 3. The reader should henceforth refer to these tables for notation.

Remark 3.5. As in [T], we use the notation $B = b_{30}b_{31} + b_{21}b_{40}$.

Having described the E_4 -term, it is now necessary to find the values of the d_4 differential on the multiplicative generators. Most of the values of d_4 follow from comparison to the classical case [T], together with a few factors of τ to balance the weights. There is only one differential that is not classical.

Lemma 3.6. $d_4(g) = h_1^4 h_4$.

Proof. By the Chow degree zero isomorphism of Theorem 2.14, we know that $h_1^4 h_4$ cannot survive the motivic May spectral sequence because $h_0^4 h_3$ is zero classically. There is only one possible differential that can kill $h_1^4 h_4$.

See also [DI] for a different proof of Lemma 3.6. $\square$

Table 3 lists the values of the d_4 differential on multiplicative generators of the E_4 -page.

3.7. The E_6 -term. We can now obtain the E_6 -page by direct computation with the d_4 differential and the Leibniz rule.

As for the E_4 -term, the multiplicative generators for the E_6 -term through the geometric 70-stem break into two groups. The first group consists of generators that survive to E_∞ and become multiplicative generators of Ext; these are listed in Table 14 as elements that occur on or before the E_6 -page. The second group consists of generators that do not survive to E_∞ ; these are listed in Table 4.

Having described the E_6 -term, it is now necessary to find the values of the d_6 differential on the multiplicative generators. Most of these values follow from comparison to the classical case [T], together with a few factors of τ to balance the weights. There are only a few differentials that are not classical.

Lemma 3.8.

- (1) $d_6(x_{56}) = h_1^2 h_5 c_0 d_0$.
- (2) $d_6(Px_{56}) = Ph_1^2 h_5 c_0 d_0$.
- (3) $d_6(B_{23}) = h_1^2 h_5 d_0 e_0$.

Proof. We have the relation $h_1x_{56} = c_0\phi$. The d_6 differential on ϕ then implies that $d_6(h_1x_{56}) = h_1^3h_5c_0d_0$, from which it follows that $d_6(x_{56}) = h_1^2h_5c_0d_0$.

The arguments for the other two differentials are similar, using the relations $h_1 \cdot Px_{56} = Pc_0 \cdot \phi$ and $h_1B_{23} = e_0\phi$. $\square$

Lemma 3.9. $d_6(c_0g^3) = h_1^{10}D_4$.

Proof. We start by computing that $h_1^2D_4$ belongs to $\langle c_0, h_4^2, h_3, h_1^3, h_1h_3 \rangle$; we will not need to worry about the indeterminacy. One can use the May d_2 differential to make this computation. All of the threefold subbrackets are strictly zero, and one of the fourfold subbrackets is also strictly zero. However, $\langle c_0, h_4^2, h_3, h_1^3 \rangle$ equals $\{0, h_1^2h_5e_0\}$.

Now shuffle to obtain that $h_1^4D_4$ is contained in $c_0\langle h_4^2, h_3, h_1^3, h_1h_3, h_1^2 \rangle$. The main point is that $h_1^4D_4$ is divisible by c_0 , which implies that there is an extension $c_0 \cdot i_1 = h_1^4D_4$ in Ext that is hidden in the May spectral sequence.

Since $h_1^6i_1 = 0$, we conclude that $h_1^{10}D_4$ must be zero in Ext . There is only one possible differential that can hit $h_1^{10}D_4$. $\square$

Remark 3.10. The value of $d_6(\Delta h_0^2Y)$ given in [T, Proposition 4.37(c)] is incorrect because it is inconsistent with machine computations of Ext_{cl} [Br1]. The value for $d_6(\Delta h_0^2Y)$ given in our table is the only possibility that is consistent with the machine computations.

Table 4 lists the values of the d_6 differential on multiplicative generators of the E_6 -page.

3.11. The E_8 -page. We can now obtain the E_8 -page by direct computation with the d_6 differential and the Leibniz rule.

Once again, the multiplicative generators for the E_8 -term through the geometric 70-stem break into two groups. The first group consists of generators that survive to E_∞ and become multiplicative generators of Ext ; these are listed in Table 14 as elements that occur on or before the E_8 -page. The second group consists of generators that do not survive to E_∞ ; these are listed in Table 5.

Once we reach the E_8 -term, we are nearly done. There are just a few more higher differentials to deal with.

Having described the E_8 -term, it is now necessary to find the values of the d_8 differential on the multiplicative generators. Once again, most of these values follow from comparison to the classical case [T], together with a few factors of τ to balance the weights. There are only a few differentials that are not classical.

Lemma 3.12.

- (1) $d_8(g^2) = h_1^8h_5$.
- (2) $d_8(w) = Ph_1^5h_5$.
- (3) $d_8(\Delta c_0g) = Ph_1^4h_5c_0$.
- (4) $d_8(Q_3) = h_1^4h_5^2$.

Proof. The Chow degree of $h_1^8h_5$ is zero. It follows from Theorem 2.14 that $h_1^8h_5$ must be zero in Ext , since $h_0^8h_4$ is zero classically. There is only one differential that can possibly hit $h_1^8h_5$.

We now know that $Ph_1^9h_5 = 0$ in Ext since $h_1^8h_5 = 0$. There is only one differential that can hit this. This shows that $d_8(w) = Ph_1^5h_5$.

Using the relation $c_0w = h_1 \cdot \Delta c_0g$, it follows that $d_8(h_1 \cdot \Delta c_0g) = Ph_1^5h_5c_0$, and then that $d_8(\Delta c_0g) = Ph_1^4h_5c_0$.

The Chow degree of $h_1^4h_5^2$ is zero. Since $h_0^4h_4^2$ is zero classically, it follows from Theorem 2.14 that $h_1^4h_5^2$ must be zero in Ext. There is only one differential that can possibly hit $h_1^4h_5^2$. $\square$

Table 5 lists the values of the d_8 differential on multiplicative generators of the E_8 -page.

3.13. The E_{12} -page. It turns out that the d_{10} differential is zero through the 70-stem, so $E_{10} = E_{12}$. The next differential to consider is d_{12} .

Lemma 3.14. *The d_{12} -differential is zero on all multiplicative generators of the E_{12} -term, except that $d_{12}(P^2Q') = Ph_0^{10}h_5i$.*

Proof. Compare to the classical case [T]. $\square$

3.15. The E_{16} -page. The d_{14} differential is zero through the 70-stem, so $E_{14} = E_{16}$.

Lemma 3.16. *The d_{16} -differential is zero on all multiplicative generators of the E_{12} -term, except that:*

- (1) $d_{16}(P^4) = h_0^{16}h_5$.
- (2) $d_{16}(\Delta^2h_4) = h_0^8h_5^2$.

Proof. Compare to the classical case [T]. $\square$

3.17. The E_{32} -page. Through the 70-stem, the last non-zero differential is d_{32} .

Lemma 3.18. *The d_{32} -differential is zero on all multiplicative generators of the E_{32} -term, except that $d_{32}(P^8) = h_0^{32}h_6$.*

Proof. Compare to the classical case [T]. $\square$

3.19. The E_∞ -term. There are no more differentials to consider in our range, and we have determined E_∞ .

The multiplicative generators for the E_∞ -term through the geometric 70-stem break into two groups, but in a different way than before. The first group consists of generators that are still multiplicative generators in Ext after hidden extensions have been considered; these are listed in Table 14. The second group consists of multiplicative generators of E_∞ that become decomposable in Ext because of a hidden extension; these are listed in Table 9.

Before ending this section, we establish that the multiplicative generator PD_4 of the E_∞ -term becomes decomposable in Ext by a hidden extension. This will be needed later in Lemma 4.12(2) to establish a hidden h_0 extension.

Lemma 3.20. $c_0 \cdot Q_2 = PD_4$.

Proof. First, compute that $h_1Q_2 = \langle h_4, h_1^2h_4, h_4, Ph_1 \rangle$ with no indeterminacy; use the May differentials $d_4(\nu_1) = h_1^2h_4^2$ and $d_4(\Delta h_1) = Ph_1h_4$. Beware that the subbracket $\langle h_1^2h_4, h_4, Ph_1 \rangle$ is not strictly zero.

Next, compute that $i_1 = \langle h_1^4, h_4, h_1^2h_4, h_4 \rangle$ with no indeterminacy; use the May differentials $d_4(g) = h_1^4h_4$ and $d_4(\nu_1) = h_1^2h_4^2$. Beware that the subbracket $\langle h_1^4, h_4, h_1^2h_4 \rangle$ is not strictly zero.

Now the shuffle

$$h_1^4 \langle h_4, h_1^2 h_4, h_4, Ph_1 \rangle = \langle h_1^4, h_4, h_1^2 h_4, h_4 \rangle Ph_1$$

implies that $h_1^5 Q_2 = Ph_1 \cdot i_1$.

Recall the hidden extension $c_0 \cdot i_1 = h_1^4 D_4$ from the proof of Lemma 3.9. This implies that $h_1^5 c_0 Q_2 = Ph_1^5 D_4$. The desired relation now follows. Note that $c_0 Q_2$ cannot equal $\tau B_{23} + PD_4$ by comparison to the classical case [Br1]. $\square$

4. HIDDEN EXTENSIONS

In order to pass from the E_∞ -term to Ext , we must resolve some hidden extensions. In this section, we deal with all possible hidden extensions by τ , h_0 , h_1 , and h_2 . We will use several different tools, including:

- (1) Classical hidden extensions [Br1].
- (2) Shuffle relations with Massey products.
- (3) Squaring operations in the sense of [Ma2].
- (4) Theorem 2.14 for hidden extensions among elements whose Chow degrees are zero.

4.1. Hidden τ extensions. By exhaustive search, the following results give all of the hidden τ extensions.

Proposition 4.2. *Table 10 lists all of the hidden τ extensions through the geometric 70-stem.*

Proof. Many of the extensions follow by comparison to the classical case. For example, there is a classical hidden extension $h_0 \cdot e_0 g = h_0^4 x$. This implies that $\tau^2 \cdot h_0 e_0 g = h_0^4 x$ motivically.

Proofs for the more subtle cases are given below. $\square$

Lemma 4.3.

- (1) $\tau \cdot \tau h_2 g^2 = Ph_1^4 h_5$.
- (2) $\tau \cdot \tau h_0 g^3 = Ph_1^4 h_5 e_0$.

Proof. Start with the relation $h_1 \cdot \tau g + h_2 f_0 = 0$, and apply the squaring operation Sq^4 . One needs that $\text{Sq}^3(\tau g) = Ph_1^2 h_5$ [Br2]. The result is the first hidden extension.

For the second, multiply the first hidden extension by e_0 . $\square$

Lemma 4.4.

- (1) $\tau \cdot B_8 = Ph_5 d_0$.
- (2) $\tau \cdot h_1^2 B_{21} = Ph_5 c_0 d_0$.
- (3) $\tau \cdot B_8 d_0 = h_0^4 X_3$.

Proof. There is a classical hidden extension $c_0 \cdot B_1 = Ph_1 h_5 d_0$ [Br1]. Motivically, there is a non-hidden relation $c_0 \cdot B_1 = h_1 B_8$. It follows that $\tau \cdot h_1 B_8 = Ph_1 h_5 d_0$ motivically.

For the second hidden extension, multiply the first hidden extension by c_0 . Note that $c_0 B_8 = h_1^2 B_{21}$ is detected in the E_∞ -term of the May spectral sequence.

For the third hidden extension, multiply the first hidden extension by d_0 , and observe that $Ph_5 d_0^2 = h_0^4 X_3$, which is detected in the E_∞ -term of the May spectral sequence. $\square$

Lemma 4.5.

- (1) $\tau \cdot Pu' = h_0^5 R_1.$
- (2) $\tau \cdot P^2 u' = h_0^9 R.$
- (3) $\tau \cdot P^3 u' = h_0^6 R'_1.$

Proof. First, use the May d_4 differential to compute that $Pu' = \langle u', h_0^3, h_0 h_3 \rangle$, with no indeterminacy. Next, use the May d_2 differential to compute that $\langle \tau, u', h_0^3 \rangle = Q'$, again with no indeterminacy.

Use a shuffle to get that $\tau \cdot Pu' = h_0 h_3 Q'$. Finally, there is a classical hidden extension $h_3 \cdot Q' = h_0^4 R_1$ [Br1], which implies that the same formula holds motivically.

The argument for the second hidden extension is similar, using the shuffle

$$\tau \cdot P^2 u' = \tau \langle u', h_0^3, h_0^5 h_4 \rangle = \langle \tau, u', h_0^3 \rangle h_0^5 h_4 = h_0^5 h_4 Q'.$$

The first equality comes from the May differential $d_8(P^2) = h_0^8 h_4$. Also, we need the classical hidden extension $h_4 \cdot Q' = h_0^4 R$ [Br1], which implies that the same formula holds motivically.

The argument for the third hidden extension is also similar, using the shuffle

$$\tau \cdot P^3 u' = \tau \langle u', h_0^3, h_0^3 i \rangle = \langle \tau, u', h_0^3 \rangle h_0^3 i = h_0^3 i Q'.$$

The first equality comes from the May differential $d_4(P^3) = h_0^6 i$. Also, we need the classical hidden extension $i \cdot Q' = h_0^3 R'_1$ [Br1]. $\square$

Lemma 4.6. $\tau \cdot k_1 = h_2 h_5 n.$

Proof. First, use the May differential $d_4(\nu) = h_0^2 h_3^2$ to compute $k = \langle d_0, h_3, h_0^2 h_3 \rangle$, with no indeterminacy. It follows from [Mi] that $\text{Sq}^0 k = \langle \text{Sq}^0 d_0, \text{Sq}^0 h_3, \text{Sq}^0 h_0^2 h_3 \rangle$, with no indeterminacy. In other words, $\text{Sq}^0 k = \langle \tau^2 d_1, h_4, \tau^2 h_1^2 h_4 \rangle$. From the classical calculation [Br2], $\text{Sq}^0 k$ also equals $\tau^3 h_2 h_5 n$.

On the other hand, the May differential $d_4(\nu_1) = h_1^2 h_4^2$ implies that $k_1 = \langle d_1, h_4, h_1^2 h_4 \rangle$, with no indeterminacy.

This shows that $\tau^4 \cdot k_1 = \tau^3 h_2 h_5 n$ in Ext, from which it follows that $\tau \cdot k_1 = h_2 h_5 n$. $\square$

Remark 4.7. In the 46-stem, $\tau \cdot u'$ does not equal $\tau^2 d_0 l$. This follows by comparison to $\text{Ext}_{A(2)}$ [I1]. Similarly, in the 49-stem, $\tau \cdot v'$ does not equal $\tau^2 e_0 l$.

4.8. Hidden h_0 extensions. By exhaustive search, the following results give all of the hidden h_0 extensions.

Proposition 4.9. *Table 11 lists all of the hidden h_0 extensions through the geometric 70-stem.*

Proof. Many of the extensions follow by comparison to the classical case. For example, there is a classical hidden extension $h_0 \cdot r = s$. This implies that $h_0 \cdot r = s$ motivically as well.

Several other extensions are implied by the hidden τ extensions established in Section 4.1. For example, the extensions $\tau \cdot Pu' = h_0^4 S_1$ and $\tau \cdot \tau h_0 d_0^2 j = h_0^5 S_1$ imply that $h_0 \cdot Pu' = \tau h_0 d_0^2 j$.

Proofs for the more subtle cases are given below. $\square$

Lemma 4.10.

- (1) $h_0 \cdot u' = \tau h_0 d_0 l.$

- (2) $h_0 \cdot v' = \tau h_0 e_0 l$.
- (3) $h_0 \cdot P v' = \tau h_0 d_0^2 k$.
- (4) $h_0 \cdot P^2 v' = \tau h_0 d_0^3 i$.

Proof. These follow by comparison to $\text{Ext}_{A(2)} [I1]$. $\square$

Lemma 4.11.

- (1) $h_0 \cdot h_2^2 g = h_1^3 h_4 c_0$.
- (2) $h_0 \cdot h_2^2 g^2 = h_1^7 h_5 c_0$.
- (3) $h_0 \cdot h_2^2 g^3 = h_1^9 D_4$.

Proof. For the first hidden extension, use the shuffle

$$h_1^3 h_4 \langle h_1, h_0, h_2^2 \rangle = \langle h_1^3 h_4, h_1, h_0 \rangle h_2^2.$$

Similarly, for the second hidden section, use the shuffle

$$h_1^7 h_5 \langle h_1, h_0, h_2^2 \rangle = \langle h_1^7 h_5, h_1, h_0 \rangle h_2^2.$$

For the third hidden extension, first recall the hidden extension $c_0 \cdot i_1 = h_1^4 D_4$ from the proof of Lemma 3.9. Use this relation to compute that

$$h_1^9 D_4 = h_1^5 i_1 \langle h_1, h_2^2, h_0 \rangle = \langle h_1^5 i_1, h_1, h_2^2 \rangle h_0.$$

Finally, compute that $h_2^2 g^3 = \langle h_1^5 i_1, h_1, h_2^2 \rangle$ from the May differential $d_4(g^3) = h_1^6 i_1$. $\square$

Lemma 4.12.

- (1) $h_0 \cdot gr = Ph_1^3 h_5 c_0$.
- (2) $h_0 \cdot lm = h_1^6 X_1$.
- (3) $h_0 \cdot m^2 = h_1^5 c_0 Q_2$.

Remark 4.13. The three parts may seem unrelated, but note that $lm = e_0 gr$ and $m^2 = g^2 r$ on the E_8 -page of the May spectral sequence.

Proof. First compute that $e_0 r = \langle \tau^2 g^2, h_2^2, h_0 \rangle$ because of the May differential $d_4(\Delta h_2 g) = \tau^2 h_2^2 g^2$. Next observe that

$$h_2 \cdot e_0 r = \langle \tau^2 g^2, h_2^2, h_0 \rangle h_2 = \langle \tau^2 g^2, h_2^2, h_0 h_2 \rangle = \langle \tau^2 h_2 g^2, h_2, h_0 h_2 \rangle.$$

None of these brackets have indeterminacy.

Use the relation $Ph_1^4 h_5 = \tau^2 h_2 g^2$ from Lemma 4.3 to write

$$h_2 \cdot e_0 r = \langle Ph_1^4 h_5, h_2, h_0 h_2 \rangle = Ph_1^3 h_5 \langle h_1, h_2, h_0 h_2 \rangle = Ph_1^3 h_5 c_0.$$

The last step is to show that $h_2 \cdot e_0 r = h_0 \cdot gr$. This follows from the calculation

$$h_0 \cdot gr = h_0 \langle h_1^3 h_4, h_1, r \rangle = \langle h_0, h_1^3 h_4, h_1 \rangle r = h_2 e_0 \cdot r.$$

This finishes the proof of part (1).

For part (2), we will prove below in Lemma 4.18 that $h_1^3 X_1 = Ph_5 c_0 e_0$. So we wish to show that $h_0 \cdot lm = Ph_1^3 h_5 c_0 e_0$. This follows immediately from part (1), using that $lm = e_0 gr$.

The proof of part (3) is similar to the proof of part (1). First, $lm = \langle \tau^2 g^3, h_2^2, h_0 \rangle$. As above, this implies that $h_2 lm = \langle \tau^2 h_2 g^3, h_2, h_0 h_2 \rangle$. Now use the (not hidden) relation $\tau^2 h_2 g^3 = h_1^6 Q_2$ to deduce that $h_2 lm = h_1^5 c_0 Q_2$. The desired formula now follows since $h_2 l = h_0 m$. $\square$

Lemma 4.14. $h_0 \cdot h_0^2 B_{22} = Ph_1 h_5 c_0 d_0$.

Proof. This follows from the hidden τ extension $\tau \cdot h_1^3 B_{21} = Ph_1 h_5 c_0 d_0$ that follows from Lemma 4.4, together with the relation $\tau h_1^3 = h_0^2 h_2$. $\square$

Remark 4.15. On the E_∞ -page, there is a relation $h_0^2 B_4 + \tau h_1 B_{21} = 0$. However, in Ext, the relation becomes $h_0^2 B_4 + \tau h_1 B_{21} = g'_2$. This follows from the analogous classical hidden relation [Br1]. Through the geometric 70-stem, this is the only example of a hidden relation of the form $h_0 \cdot x + h_1 \cdot y$, $h_0 \cdot x + h_2 \cdot y$, or $h_1 \cdot x + h_2 \cdot y$.

4.16. Hidden h_1 extensions. By exhaustive search, the following results give all of the hidden h_1 extensions.

Proposition 4.17. *Table 12 lists all of the hidden h_1 extensions through the geometric 70-stem.*

Proof. Many of the extensions follow by comparison to the classical case. For example, there is a classical hidden extension $h_1 \cdot x = h_2^2 d_1$. This implies that there is a motivic hidden extension $h_1 \cdot x = \tau h_2^2 d_1$.

Proofs for the more subtle cases are given below. $\square$

Lemma 4.18.

- (1) $h_1 \cdot \tau h_1 G = h_5 c_0 e_0$.
- (2) $h_1 \cdot h_1 B_3 = h_5 d_0 e_0$.
- (3) $h_1 \cdot \tau P h_1 G = Ph_5 c_0 e_0$.
- (4) $h_1 \cdot h_1^2 X_3 = h_5 c_0 d_0 e_0$.

Proof. The element $\tau h_1 G$ is represented in the May spectral sequence by $\tau \Delta_1 h_1^3$, which equals $\Delta_1 h_0^2 h_2$ on the May E_4 -page. We can then compute that $\tau h_1 G = \langle h_5, h_2 g, h_0^2 \rangle$ using the May differential $d_4(\Delta_1 h_2) = h_2 h_5 g$. Next shuffle to obtain $h_1 \cdot \tau h_1 G = \langle h_5, h_2 g, h_0^2 \rangle h_1 = h_5 \langle h_2 g, h_0^2, h_1 \rangle$. Finally, compute that $c_0 e_0 = \langle h_2 g, h_0^2, h_1 \rangle$ with the May differential $d_2(h_0(1)^2 b_{21}) = h_2^2 h_0(1)^2$. This establishes the first hidden extension.

For the second hidden extension, begin by computing that $h_1 h_5 d_0 e_0 = \langle h_5 c_0 e_0, h_0, h_2^2 \rangle$ using the May differential $d_2(h_0(1)) = h_0 h_2^2$. From part (1), this equals $\langle \tau h_1^2 G, h_0, h_2^2 \rangle$, which equals $h_1^2 \langle \tau G, h_0, h_2^2 \rangle$ because there is no indeterminacy. This shows that $h_5 d_0 e_0$ is divisible by h_1 . The only possible hidden extension is $h_1 \cdot h_1 B_3 = h_5 d_0 e_0$.

For the third hidden extension, start with the relation $h_1 \cdot \tau P G = Ph_1 \cdot \tau G$ because there is no possible hidden relation. Therefore, using part (1),

$$h_1^3 \cdot \tau P G = Ph_1 \cdot h_1^2 \cdot \tau G = Ph_1 h_5 c_0 e_0.$$

It follows that $h_1 \cdot \tau P h_1 G = Ph_5 c_0 e_0$.

For the fourth hidden extension, use part (1) to conclude that $h_5 c_0 d_0 e_0$ is divisible by h_1 . The only possibility is that $h_1 \cdot h_1^2 X_3 = h_5 c_0 d_0 e_0$. $\square$

Lemma 4.19. $h_1 \cdot h_1^2 B_6 = \tau h_2^2 d_1 g$.

Proof. Begin with the hidden extension $h_1 \cdot x = \tau h_2^2 d_1$, which follows by comparison to the classical case [Br1]. Next compute that $\tau h_2^2 d_1 g = \langle \tau h_2^2 d_1, h_1^2, h_1^2 h_4 \rangle$, using the May differential $d_4(g) = h_1^4 h_4$. It follows that $\tau h_2^2 d_1 g = \langle h_1 x, h_1^2, h_1^2 h_4 \rangle$, which equals $h_1 \langle x, h_1^2, h_1^2 h_4 \rangle$ because there is no indeterminacy. Therefore, $\tau h_2^2 d_1 g$ is divisible by h_1 , and the only possibility is that $h_1 \cdot h_1^2 B_6 = \tau h_2^2 d_1 g$. $\square$

Lemma 4.20. $h_1 \cdot h_1 D_{11} = \tau^2 c_1 g^2$.

Proof. Begin by computing that $h_1 D_{11} = \langle y, h_1^2, h_1^2 h_4 \rangle$, using the May differential $d_4(g) = h_1^4 h_4$ and the relation $\Delta h_3^2 g = \Delta h_1^2 d_1$. Also recall the hidden extension $h_1 \cdot y = \tau^2 c_1 g$, which follows by comparison to the classical case [Br1].

It follows that

$$h_1^2 D_{11} = \langle h_1 y, h_1^4, h_4 \rangle = \langle \tau^2 c_1 g, h_1^4, h_4 \rangle$$

because there is no indeterminacy. Finally, use the May differential $d_4(g) = h_1^4 h_4$ again to compute that $\tau^2 c_1 g^2 = \langle \tau^2 c_1 g, h_1^2, h_1^2 h_4 \rangle$. $\square$

Lemma 4.21. $h_1 \cdot C_0$ equals zero, not $h_0 h_5 l$.

Proof. First compute that C_0 belongs to $\langle h_0 h_3^2, h_0, h_1, \tau h_1 g_2 \rangle$ using the May differentials $d_4(\nu) = h_0^2 h_3^2$ and $d_4(x_{47}) = \tau h_1^2 g_2$. Note that the subbracket $\langle h_0 h_3^2, h_0, h_1 \rangle$ is strictly zero, but the subbracket $\langle h_0, h_1, \tau h_1 g_2 \rangle$ equals $\{0, h_0 h_2 g_2\}$. We will not need to compute the indeterminacy in the fourfold bracket.

Now a shuffle implies that $h_1 \cdot C_0$ belongs to $\langle h_1, h_0 h_3^2, h_0, h_1 \rangle \tau h_1 g_2$. For degree reasons, the bracket $\langle h_1, h_0 h_3^2, h_0, h_1 \rangle$ consists of elements spanned by f_0 and $\tau h_1^3 h_4$. But the products $h_1 \cdot f_0$ and $h_1 \cdot \tau h_1^3 h_4$ are both zero, so $\langle h_1, h_0 h_3^2, h_0, h_1 \rangle \tau h_1 g_2$ must be zero. Therefore, $h_1 \cdot C_0$ is zero. $\square$

Lemma 4.22. $h_1 \cdot r_1 = s_1$.

Proof. This follows immediately from Theorem 2.14 and the classical relation $h_0 \cdot r = s$. $\square$

Lemma 4.23. $h_1 \cdot h_1^2 q_1 = h_0^4 X_3$.

Proof. Apply Sq^6 to the relation $h_2 r = h_1 q$ to obtain that $h_3 r^2 = h_1^2 \text{Sq}^5(q)$. Next, observe that $\text{Sq}^5(q) = h_1 q_1$ by comparison to the classical case [Br2].

By comparison to the classical case, there is a relation $h_3 r = h_0^2 x + \tau h_2^2 n$, so $h_3 r^2 = h_0^2 r x$. Finally, use the hidden extension $h_0 \cdot r = s$ and the non-hidden relation $s x = h_0^3 X_3$. $\square$

4.24. Hidden h_2 extensions. By exhaustive search, the following results give all of the hidden h_2 extensions.

Proposition 4.25. Table 13 lists all of the hidden h_2 extensions through the geometric 70-stem.

Proof. Many of the extensions follow by comparison to the classical case. For example, there is a classical hidden extension $h_2 \cdot Q_2 = h_5 k$. This implies that the same formula holds motivically.

Also, many extensions are implied by hidden h_0 extensions that we already established in Section 4.8. For example, There is a hidden extension $h_0 \cdot h_2^2 g = h_1^3 h_4 c_0$. This implies that there is also a hidden extension $h_2 \cdot h_0 h_2 g = h_1^3 h_4 c_0$.

Proofs for the more subtle cases are given below. $\square$

Remark 4.26. We established the extensions

- (1) $h_2 \cdot e_0 r = P h_1^3 h_5 c_0$
- (2) $h_2 \cdot l m = h_1^5 c_0 Q_2$

in the proof of Lemma 4.12. The extension $h_2 \cdot k m = h_1^6 X_1$ follows from Lemma 4.12 and the relation $h_2 k = h_0 l$.

Lemma 4.27. $h_2 \cdot h_2 B_2 = h_1 h_5 c_0 d_0$.

Proof. First compute that $h_2 B_2 = \langle g_2, h_0^3, h_2^2 \rangle$, with no indeterminacy, using the May differential $d_6(Y) = h_0^3 g_2$. Then $h_2 \cdot h_2 B_2$ equals $\langle g_2, h_0^3, h_2^3 \rangle$, because there is no indeterminacy. This bracket equals $\langle g_2, h_0^3, h_1^2 h_3 \rangle$, which equals $\langle g_2, h_0^3, h_1 \rangle h_1 h_3$ since there is no indeterminacy. Compute that the bracket $\langle g_2, h_0^3, h_1 \rangle$ equals B_1 , using the May differential $d_6(Y) = h_0^3 g_2$.

We have now shown that $h_2 \cdot h_2 B_2$ equals $h_1 h_3 \cdot B_1$. It remains to show that there is a hidden extension $h_3 \cdot B_1 = h_5 c_0 d_0$. First observe that $B_1 \cdot h_1^2 d_0 = h_1^3 B_{21}$ by a non-hidden relation. This implies that $B_1 \cdot \tau h_1^2 d_0 = Ph_1 h_5 c_0 d_0$ by Lemma 4.4.

Now there is a hidden extension $h_3 \cdot Ph_1 = \tau h_1^2 d_0$, so $B_1 \cdot h_3 \cdot Ph_1 = Ph_1 h_5 c_0 d_0$. The only possibility is that $h_3 \cdot B_1 = h_5 c_0 d_0$. $\square$

Lemma 4.28. $h_2 \cdot B_6 = \tau e_1 g$.

Proof. We will first prove that

$$\langle \tau, B_6, h_1^2 h_3 \rangle = h_2 C_0,$$

with no indeterminacy. The May differentials $d_2(b_{30} b_{40} h_1(1)) = \tau B_6$ and $d_2(B h_1 b_{21} h_1(1)) = h_1^2 h_3 B_6$ imply that the bracket equals $\tau B h_1 b_{21} h_1(1) + h_1^2 h_3 b_{30} b_{40} h_1(1)$. But this expression equals $h_2 C_0$ on the May E_4 -page because of the differential

$$d_2(b_{21} b_{30} b_{40} h_1(1)) = h_2^3 b_{30} b_{40} h_1(1) + \tau B h_1 b_{21} h_1(1) + h_1^3 b_{30} b_{22} b_{40}.$$

This means that

$$\langle \tau, B_6, h_2^3 \rangle = h_2 C_0.$$

If $h_2 \cdot B_6$ were zero, then this would imply that

$$\langle \tau, B_6, h_2 \rangle h_2^2 = h_2 C_0.$$

However, C_0 cannot be divisible by h_2 . $\square$

5. h_1 -LOCAL COHOMOLOGY OF MOTIVIC $A(3)$

Because h_1^k is non-zero in Ext , it makes sense to invert h_1 and compute $\text{Ext}[h_1^{-1}]$. In fact, $\text{Ext}[h_1^{-1}]$ is practically computable in a much larger range than Ext . For our purposes, it is enough to consider $\text{Ext}_{A(3)}[h_1^{-1}]$ because this simpler calculation captures everything that we need to know in the range under study. Here $A(3)$ is the $\mathbb{M}_2$ -subalgebra of the Steenrod algebra A generated by Sq^1 , Sq^2 , Sq^4 , and Sq^8 .

Since localization is exact, we can invert h_1 in the E_2 -term of the motivic May spectral sequence for $A(3)$; then we have a spectral sequence that converges to $\text{Ext}_{A(3)}[h_1^{-1}]$.

When we do this, we find that some of the usual generators can be written in terms of other generators. For example, the relation $h_0(1)^2 = b_{20} b_{21} + h_1^2 b_{30}$ implies that $b_{30} = h_1^{-2} h_0(1)^2 + h_1^{-2} b_{20} b_{21}$. Similarly, some relations simplify, and others become redundant. For example, the relation $h_0 h_1 = 0$ implies that $h_0 = 0$.

Proposition 5.1. *The h_1 -local E_2 -term of the motivic May spectral sequence for $A(3)$ is the $\mathbb{M}_2[h_1^{\pm 1}]$ -algebra generated by the elements in the following table, subject to the single relation $h_3 h_0(1) = 0$. The values of the May d_2 differential are also given in the table.*

generator	(m, s, f, w)	d_2
h_3	$(1, 7, 1, 4)$	τh_1^3 $h_1^2 h_3$
b_{20}	$(4, 4, 2, 2)$	
$h_0(1)$	$(4, 7, 2, 4)$	
b_{21}	$(4, 10, 2, 6)$	
b_{31}	$(6, 26, 2, 14)$	
b'_{40}	$(10, 30, 4, 16)$	

The element b'_{40} is defined to be $h_1^2 b_{40} + b_{20} b_{31}$.

Proof. This follows immediately by comparison to the motivic May spectral sequence for A . $\square$

Proposition 5.2. *The h_1 -local E_∞ -term of the motivic May spectral sequence for $A(3)$ is the $\frac{\mathbb{M}_2}{\tau}[h_1^{\pm 1}]$ -algebra generated by the elements in the following table, subject to the single relation $h_1^2 e_0^2 + c_0^2 g = 0$.*

generator	(m, s, f, w)	E_2 description
P	$(8, 8, 4, 4)$	b_{20}^2
c_0	$(5, 8, 3, 5)$	$h_1 h_0(1)$
e_0	$(8, 17, 4, 10)$	$b_{21} h_0(1)$
g	$(8, 20, 4, 12)$	b_{21}^2
b_{31}	$(6, 26, 2, 14)$	
b'_{40}	$(10, 30, 4, 16)$	

Proof. After the d_2 differential, there are no more differentials, and $E_3 = E_\infty$. $\square$

At this point, we almost completely know $\text{Ext}_{A(3)}[h_1^{-1}]$. There is just one possible hidden extension.

Lemma 5.3. $h_1^2 e_0^2 + c_0^2 g = h_1^4 b'_{40}$ in $\text{Ext}_{A(3)}[h_1^{-1}]$,

Remark 5.4. The formula has been verified by machine computation. Unfortunately, we have been unable to find a conceptual proof of Lemma 5.3. Use of brackets and squaring operations has not yielded any results.

Remark 5.5. This obscure algebraic hidden extension has significant consequences for classical Adams differentials, as shown in [12].

Theorem 5.6. $\text{Ext}_{A(3)}[h_1^{-1}]$ is equal to the free polynomial $\frac{\mathbb{M}_2}{\tau}[h_1^{\pm 1}]$ -algebra generated by the elements in the following table.

generator	(s, f, w)	E_2 description
P	$(8, 4, 4)$	b_{20}^2
c_0	$(8, 3, 5)$	$h_1 h_0(1)$
e_0	$(17, 4, 10)$	$b_{21} h_0(1)$
g	$(20, 4, 12)$	b_{21}^2
b_{31}	$(26, 2, 14)$	

Proof. This follows immediately from Proposition 5.2 and Lemma 5.3. $\square$

Corollary 5.7. *Through the geometric 70-stem, the h_1 -local elements of Ext are precisely the elements indicated in the charts with red arrows of slope 1.*

Proof. Each of these elements is detected by the map $\text{Ext}[h_1^{-1}] \rightarrow \text{Ext}_{A(3)}[h_1^{-1}]$. $\square$

We give one specific consequence of these h_1 -local calculations. The following corollary is the algebraic key to a number of Adams differentials beyond the 50-stem.

Corollary 5.8. *In Ext , there is a hidden extension $e_0^3 + d_0 \cdot e_0 g = h_1^5 B_1$.*

Proof. This extension can be detected by the map $\text{Ext} \rightarrow \text{Ext}_{A(3)}[h_1^{-1}]$. Under the induced map on E_∞ -pages of the respective May spectral sequences, we have that d_0 goes to $h_1^{-2} c_0^2$; e_0 goes to e_0 ; and $e_0 g$ goes to $e_0 g$.

The map $\text{Ext} \rightarrow \text{Ext}_{A(3)}[h_1^{-1}]$ then takes d_0 to $h_1^{-2} c_0^2$; and e_0 to e_0 . However, for degree reasons, it is possible that $e_0 g$ maps to either $e_0 g$ or $e_0 g + h_1^3 c_0 b_{31}$. (In fact, it maps to $e_0 g + h_1^3 c_0 b_{31}$, but we will not need this.)

In either case, $e_0^3 + d_0 \cdot e_0 g$ maps to a non-zero element in $\text{Ext}_{A(3)}[h_1^{-1}]$. It follows that there is a hidden extension, and there is only one possibility. $\square$

6. TABLES

Table 1: May E_2 -term generators in stems 0–70

	(m, s, f, w)	d_2	description
h_0	(1, 0, 1, 0)		h_{10}
h_1	(1, 1, 1, 1)		h_{11}
h_2	(1, 3, 1, 2)		h_{12}
b_{20}	(4, 4, 2, 2)	$\tau h_1^3 + h_0^2 h_2$	h_{20}^2
h_3	(1, 7, 1, 4)		h_{13}
$h_0(1)$	(4, 7, 2, 4)	$h_0 h_2^2$	$h_{20} h_{21} + h_{11} h_{30}$
b_{21}	(4, 10, 2, 6)	$h_2^3 + h_1^2 h_3$	h_{21}^2
b_{30}	(6, 12, 2, 6)	$\tau h_1 b_{21} + h_3 b_{20}$	h_{30}^2
h_4	(1, 15, 1, 8)		h_{14}
$h_1(1)$	(4, 16, 2, 9)	$h_1 h_3^2$	$h_{21} h_{22} + h_{12} h_{31}$
b_{22}	(4, 22, 2, 12)	$h_3^3 + h_2^2 h_4$	h_{22}^2
b_{31}	(6, 26, 2, 14)	$h_4 b_{21} + h_2 b_{22}$	h_{31}^2
b_{40}	(8, 28, 2, 14)	$h_4 b_{30} + \tau h_1 b_{31}$	h_{40}^2
h_5	(1, 31, 1, 16)		h_{15}
$h_2(1)$	(4, 34, 2, 18)	$h_2 h_4^2$	$h_{22} h_{23} + h_{13} h_{32}$
$h_0(1, 3)$	(7, 38, 3, 20)	$h_4^2 h_0(1) + h_0 h_2 h_2(1)$	$h_{50} h_{11} h_{13} + h_{40} h_{11} h_{23} +$ $+ h_{20} h_{41} h_{13} + h_{20} h_{31} h_{23}$
b_{23}	(4, 46, 2, 24)	$h_4^3 + h_3^2 h_5$	h_{23}^2
$h_0(1, 2)$	(9, 46, 3, 24)	$h_3 h_0(1, 3)$	$h_{30} h_{31} h_{32} + h_{30} h_{41} h_{22} +$ $+ h_{40} h_{21} h_{32} + h_{40} h_{41} h_{12} +$ $+ h_{50} h_{21} h_{22} + h_{50} h_{31} h_{12}$
b_{32}	(6, 54, 2, 28)	$h_5 b_{22} + h_3 b_{23}$	h_{32}^2
b_{41}	(8, 58, 2, 30)	$h_5 b_{31} + h_2 b_{32}$	h_{41}^2
b_{50}	(10, 60, 2, 30)	$h_5 b_{40} + \tau h_1 b_{41}$	h_{50}^2
h_6	(1, 63, 1, 32)		h_{16}
$h_3(1)$	(4, 70, 2, 36)	$h_3 h_5^2$	$h_{23} h_{24} + h_{14} h_{33}$

Table 2: May E_2 -term relations in stems 0–46

relation	(m, s, f, w)
$h_0 h_1$	(2, 1, 2, 1)
$h_1 h_2$	(2, 4, 2, 3)
$h_2 b_{20} = h_0 h_0(1)$	(5, 7, 3, 4)
$h_2 h_3$	(2, 10, 2, 6)
$h_2 h_0(1) = h_0 b_{21}$	(5, 10, 3, 6)
$h_3 h_0(1)$	(5, 14, 3, 8)
$h_0(1)^2 = b_{20} b_{21} + h_1^2 b_{30}$	(8, 14, 4, 8)
$h_0 h_1(1)$	(5, 16, 3, 9)
$h_3 b_{21} = h_1 h_1(1)$	(5, 17, 3, 10)
$b_{20} h_1(1) = h_1 h_3 b_{30}$	(8, 20, 4, 11)
$h_3 h_4$	(2, 22, 2, 12)
$h_3 h_1(1) = h_1 b_{22}$	(5, 23, 3, 13)
$h_0(1) h_1(1)$	(8, 23, 4, 13)
$b_{20} b_{22} = h_0^2 b_{31} + h_3^2 b_{30}$	(8, 26, 4, 14)
$b_{22} h_0(1) = h_0 h_2 b_{31}$	(8, 29, 4, 16)
$h_4 h_1(1)$	(5, 31, 3, 17)
$h_1(1)^2 = b_{21} b_{22} + h_2^2 b_{31}$	(8, 32, 4, 18)
$h_1 h_2(1)$	(5, 35, 3, 19)
$h_4 b_{22} = h_2 h_2(1)$	(5, 37, 3, 20)
$b_{20} h_2(1) = h_0 h_0(1, 3)$	(8, 38, 4, 20)
$h_2 h_0(1, 3) = h_0 h_4 b_{31}$	(8, 41, 4, 22)
$h_0(1) h_2(1) = h_0 h_4 b_{31}$	(8, 41, 4, 22)
$b_{21} h_2(1) = h_2 h_4 b_{31}$	(8, 44, 4, 24)
$h_0(1) h_0(1, 3) = h_1^2 h_4 b_{40} + h_4 b_{20} b_{31}$	(11, 45, 5, 24)
$h_4 h_5$	(2, 46, 2, 24)
$b_{30} h_2(1) = h_0 h_0(1, 2) + h_2 h_4 b_{40}$	(10, 46, 4, 24)

Table 2: May E_2 -term relations in stems 48–70	
relation	(m, s, f, w)
$b_{21}h_0(1, 3) = h_1^2h_0(1, 2) + h_4b_{31}h_0(1)$	(11, 48, 5, 26)
$h_4h_2(1) = h_2b_{23}$	(5, 49, 3, 26)
$h_1(1)h_2(1)$	(8, 50, 4, 27)
$b_{30}h_0(1, 3) = b_{20}h_0(1, 2) + h_4b_{40}h_0(1)$	(13, 50, 5, 26)
$b_{23}h_0(1) = h_4h_0(1, 3)$	(8, 53, 4, 28)
$h_0(1)h_0(1, 2) = h_4b_{40}b_{21} + h_4b_{30}b_{31}$	(13, 53, 5, 28)
$h_1(1)h_0(1, 3) = h_1h_3h_0(1, 2)$	(11, 54, 5, 29)
$b_{21}b_{23} = h_1^2b_{32} + h_4^2b_{31}$	(8, 56, 4, 30)
$b_{30}b_{23} = b_{20}b_{32} + h_0^2b_{41} + h_4^2b_{40}$	(10, 58, 4, 30)
$b_{22}h_0(1, 3) = h_3^2h_0(1, 2) + h_0b_{31}h_2(1)$	(11, 60, 5, 32)
$b_{32}h_0(1) = h_4h_0(1, 2) + h_0h_2b_{41}$	(10, 61, 4, 32)
$b_{23}h_1(1) = h_1h_3b_{32}$	(8, 62, 4, 33)
$h_5h_2(1)$	(5, 65, 3, 34)
$b_{22}b_{23} = h_2(1)^2 + h_3^2b_{32}$	(8, 68, 4, 36)
$h_5h_0(1, 3)$	(8, 69, 4, 36)

Table 3: Temporary generators of the May E_4 -term			
	(m, s, f, w)	description	d_4
P	(8, 8, 4, 4)	b_{20}^2	$h_0^4h_3$
ν	(7, 15, 3, 8)	h_2b_{30}	$h_0^2h_3^2$
g	(8, 20, 4, 12)	b_{21}^2	$h_1^4h_4$
Δ	(12, 24, 4, 12)	b_{30}^2	$\tau^2h_2g + Ph_4$
ν_1	(7, 33, 3, 18)	h_3b_{31}	$h_1^2h_4^2$
x_{34}	(7, 34, 5, 18)	$h_0^3h_2(1) + h_0h_4^2b_{20}$	
x_{35}	(10, 35, 4, 18)	$h_0h_3b_{40}$	x_{34}
Y	(16, 45, 6, 24)	$Bh_0(1)$	
x_{47}	(13, 47, 5, 25)	$h_2b_{40}h_1(1)$	$\tau h_1^2g_2$
x_{49}	(10, 49, 4, 26)	$h_2h_0(1, 2)$	$h_0h_3c_2$
ϕ	(17, 49, 7, 27)	Bh_1b_{21}	
Δ_1	(12, 52, 4, 28)	b_{31}^2	$h_5g + h_3g_2$
X	(23, 54, 9, 28)	$Bh_0b_{20}b_{30}$	
x_{56}	(21, 56, 9, 31)	$Bh_1b_{21}h_0(1)$	
Γ	(16, 56, 4, 28)	b_{40}^2	$\Delta h_5 + \tau^2\Delta_1h_2$
x'_{56}	(24, 56, 10, 30)	PBb_{21}	
ϕ'	(23, 57, 9, 30)	$Bh_0b_{30}h_0(1)$	
x_{59}	(19, 59, 7, 31)	$h_2b_{30}b_{40}h_1(1)$	τ^2e_1g
x_{63}	(20, 63, 8, 33)	$h_1b_{20}b_{30}h_0(1, 2) +$ $+\tau h_1^2b_{31}b_{40}h_0(1) +$ $+\tau b_{20}b_{31}^2h_0(1)$	$\tau h_5d_0e_0$
x_{65}	(20, 65, 8, 34)	$h_0h_3b_{20}b_{31}b_{40}$	h_0^3A''
x_{68}	(16, 68, 6, 36)	$h_3^2b_{31}b_{40} + \tau b_{31}^2h_1(1)$	τs_1
ν_2	(7, 69, 3, 36)	h_4b_{32}	$h_2^2h_5^2$
x_{69}	(13, 69, 5, 36)	$h_3b_{40}h_2(1)$	$h_0^2d_2$

Table 4: Temporary generators of the May E_6 -term

	(m, s, f, w)	description	d_6
P^2	(16, 16, 8, 8)		
Δh_3	(13, 31, 5, 16)		
Pi	(23, 31, 11, 16)	$P^2\nu$	$h_0^5 s$
Pr	(22, 38, 10, 20)	$P\Delta h_2^2$	$h_0^6 x$
g^2	(16, 40, 8, 24)		
Y	(16, 45, 6, 24)		$h_0^3 g_2$
w	(21, 45, 9, 25)	$\Delta h_1 g$	
Δ^2	(24, 48, 8, 24)		
ϕ	(17, 49, 7, 27)		$h_1^3 h_5 d_0$
$\Delta c_0 g$	(25, 52, 11, 29)		
X	(23, 54, 9, 28)		$Ph_0 h_5 d_0$
PQ'	(37, 55, 17, 28)	$P^2 \Delta h_0^2 \nu$	$h_0^9 X$
x_{56}	(21, 56, 9, 31)		$h_1^2 h_5 c_0 d_0$
x'_{56}	(24, 56, 10, 30)		$Ph_1^2 h_5 d_0$
ϕ'	(23, 57, 9, 30)		$Ph_0 h_5 e_0$
Px_{56}	(29, 64, 13, 35)		$Ph_1^2 h_5 c_0 d_0$
Px'_{56}	(32, 64, 14, 34)		$P^2 h_1^2 h_5 d_0$
B_{23}	(24, 65, 10, 36)	Yg	$h_1^2 h_5 d_0 e_0$
$P\phi'$	(31, 65, 13, 34)		$P^2 h_0 h_5 e_0$
Q_3	(13, 67, 5, 36)	$\Delta_1 h_4$	
$c_0 g^3$	(29, 68, 15, 41)		$h_1^{10} D_4$
$\Delta h_0^2 Y$	(30, 69, 12, 36)		$\Delta h_0^5 g_2 + h_0 h_5 d_0 i$
$\Gamma h_0 h_3^2$	(19, 70, 7, 36)		

Table 5: Temporary generators for the May E_8 -term

element	(m, s, f, w)	d_8
P^2	(16, 16, 8, 8)	$h_0^8 h_4$
Δh_3	(13, 31, 5, 16)	$h_0^4 h_4^2$
g^2	(16, 40, 8, 24)	$h_1^8 h_5$
w	(21, 45, 9, 25)	$Ph_1^5 h_5$
Δ^2	(24, 48, 8, 24)	$P^2 h_5$
$\Delta c_0 g$	(25, 52, 11, 29)	$Ph_1^4 h_5 c_0$
Q_3	(13, 67, 5, 36)	$h_1^4 h_5^2$
$\Gamma h_0 h_3^2$	(19, 70, 7, 36)	$h_0^4 p'$

Table 6: Temporary generators for the May E_{12} -term

element	(m, s, f, w)	d_{12}
P^4	(32, 32, 16, 16)	
$P^2 Q'$	(45, 63, 21, 32)	$Ph_0^{10} h_5 i$
$\Delta^2 h_4$	(25, 63, 9, 32)	

Table 7: Temporary generators for the May E_{16} -term

	(m, s, f, w)	d_{16}
P^4	(32, 32, 16, 16)	$h_0^{16} h_5$
$\Delta^2 h_4$	(25, 63, 9, 32)	$h_0^8 h_5^2$

Table 8: Temporary generators for the May E_{32} -term

element	(m, s, f, w)	d_{32}
P^8	(64, 64, 32, 32)	$h_0^{32} h_6$

Table 9: Temporary generators for the May E_∞ -term

element	(m, s, f, w)	description	occurs
s	(13, 30, 7, 16)	$Ph_4h_0(1) + h_0^3b_{20}b_{31}$	4
P^2s	(29, 46, 15, 24)		12
S_1	(25, 54, 11, 28)	h_0^2X	8
g'_2	(23, 60, 11, 32)	$P\Delta_1h_0^3$	6
τPG	(22, 62, 10, 33)	$\tau P\Delta_1h_1^2$	6
Ph_5i	(24, 62, 12, 32)		8
P^4s	(45, 62, 23, 32)		32
PD_4	(22, 65, 10, 35)		6
s_1	(13, 67, 7, 37)	$h_5b_{21}^2h_1(1) + h_1^3b_{21}b_{32}$	4
Ph_5^2	(10, 70, 6, 36)		6
τP^2G	(30, 70, 14, 37)		12
P^2S_1	(41, 70, 19, 36)		12

Table 10: Hidden τ extensions in stems 0–54

(s, f, w)	x	$\tau \cdot x$
(30, 11, 16)	Pc_0d_0	h_0^4s
(37, 9, 20)	τh_0e_0g	h_0^4x
(37, 10, 20)	$\tau h_0^2e_0g$	h_0^5x
(41, 5, 22)	h_1f_1	$h_0^2c_2$
(43, 9, 24)	τh_2g^2	$Ph_1^4h_5$
(46, 19, 24)	$P^3c_0d_0$	$P^2h_0^4s$
(53, 9, 28)	B_8	Ph_5d_0
(53, 17, 28)	$\tau Ph_0d_0^2e_0$	h_0^7x'
(53, 18, 28)	$\tau Ph_0^2d_0^2e_0$	h_0^8x'
(54, 10, 29)	h_1B_8	$Ph_1h_5d_0$
(54, 15, 28)	Pu'	$h_0^4S_1$
(54, 16, 28)	$\tau h_0d_0^2j$	$h_0^5S_1$
(54, 17, 28)	$\tau^2Ph_1d_0^2e_0$	$h_0^6S_1$

Table 10: Hidden τ extensions in stems 58–70

(s, f, w)	x	$\tau \cdot x$
(58, 8, 30)	h_1Q_2	$h_0^2D_2$
(60, 13, 34)	τh_0g^3	$Ph_1^4h_5e_0$
(61, 12, 33)	$h_1^2B_{21}$	$Ph_5c_0d_0$
(61, 13, 32)	$x'c_0$	$h_0^4X_1$
(62, 13, 34)	$h_1^3B_{21}$	$Ph_1h_5c_0d_0$
(62, 19, 32)	P^2u'	$Ph_0^7h_5i$
(62, 20, 32)	$\tau Ph_0d_0^2j$	$Ph_0^8h_5i$
(62, 21, 32)	$\tau^2P^2h_1d_0^2e_0$	$Ph_0^9h_5i$
(62, 27, 32)	$P^5c_0d_0$	$P^4h_0^4s$
(64, 8, 34)	h_1X_2	h_0^2A''
(65, 7, 35)	k_1	h_2h_5n
(65, 9, 34)	$h_1h_3Q_2$	$h_0^2h_3D_2$
(66, 12, 34)	$h_1^2q_1$	$h_0^2D'_2$
(67, 13, 36)	B_8d_0	$h_0^4X_3$
(68, 12, 36)	$\tau h_0h_2B_{23}$	$h_0^4G_{21} + h_5d_0i$
(69, 7, 36)	$\tau h_1^2Q_3$	h_0^3p'
(69, 25, 36)	$\tau P^3h_0d_0^2e_0$	$P^2h_0^7x'$
(69, 26, 36)	$\tau P^3h_0^2d_0^2e_0$	$P^2h_0^8x'$
(70, 6, 36)	τh_2Q_3	Ph_5^2
(70, 7, 36)	$\tau h_0h_2Q_3$	$Ph_0h_5^2$
(70, 23, 36)	P^3u'	$P^2h_0^4S_1$
(70, 24, 36)	$\tau P^2h_0d_0^2j$	$h_0^5S_1$
(70, 25, 36)	$\tau P^2h_0^2d_0^2j$	$h_0^6S_1$

Table 11: Hidden h_0 extensions

(s, f, w)	x	$h_0 \cdot x$
(26, 7, 16)	$h_2^2 g$	$h_1^3 h_4 c_0$
(30, 7, 16)	r	s
(46, 11, 28)	$h_2^2 g^2$	$h_1^7 h_5 c_0$
(46, 12, 25)	u'	$\tau h_0 d_0 l$
(46, 15, 24)	i^2	$P^2 s$
(49, 12, 27)	v'	$\tau h_0 e_0 l$
(50, 11, 28)	gr	$Ph_1^3 h_5 c_0$
(54, 16, 29)	Pu'	$\tau h_0 d_0^2 j$
(54, 11, 28)	R_1	S_1
(56, 11, 29)	Q_1	$\tau h_2 x'$
(57, 16, 31)	Pv'	$\tau h_0 d_0 e_0 j$
(60, 8, 32)	B_3	$h_5 k$
(60, 16, 33)	$d_0 u'$	$\tau h_0 e_0^2 j$
(62, 20, 33)	$P^2 u'$	$\tau Ph_0 d_0^2 j$
(62, 12, 32)	$h_0 R$	$Ph_5 i$
(62, 13, 34)	$h_0^2 B_{22}$	$Ph_1 h_5 c_0 d_0$
(62, 23, 32)	$P^2 i^2$	$P^4 s$
(63, 8, 34)	X_2	$h_5 l$
(63, 15, 38)	$h_0 h_2 g^3$	$h_1^7 h_5 c_0 e_0$
(63, 16, 35)	$e_0 u'$	$\tau h_0 e_0^2 k$
(64, 8, 34)	$h_2 A'$	$\tau^2 d_1^2$
(64, 15, 34)	U	$Ph_2 x'$
(65, 20, 35)	$P^2 v'$	$\tau h_0 d_0^3 i$
(66, 15, 40)	$h_2^2 g^3$	$h_1^9 D_4$
(66, 16, 37)	$e_0 v'$	$\tau h_0 e_0^2 l$
(67, 14, 38)	lm	$h_1^6 X_1$
(68, 15, 36)	$h_0 G_{11}$	$\tau h_1 d_0 x'$
(68, 20, 37)	$Pd_0 u'$	$\tau h_0 d_0^3 j$

Table 11: Hidden h_0 extensions

(s, f, w)	x	$h_0 \cdot x$
(70, 24, 37)	$P^3 u'$	$\tau P^2 h_0 d_0^2 j$
(70, 15, 40)	m^2	$h_1^5 c_0 Q_2$
(70, 19, 36)	$h_0 R_1'$	$P^2 S_1$

Table 12: Hidden h_1 extensions

(s, f, w)	x	$h_1 \cdot x$
(38, 6, 21)	x	$\tau h_2^2 d_1$
(39, 7, 21)	y	$\tau^2 c_1 g$
(56, 8, 31)	$\tau h_1 G$	$h_5 c_0 e_0$
(58, 10, 33)	$h_1^2 B_6$	$\tau h_2^2 d_1 g$
(59, 11, 33)	$h_1 D_{11}$	$\tau^2 c_1 g^2$
(62, 9, 34)	$h_1 B_3$	$h_5 d_0 e_0$
(62, 10, 33)	X_1	τPG
(64, 8, 35)	C'	τd_1^2
(64, 12, 35)	$\tau Ph_1 G$	$Ph_5 c_0 e_0$
(67, 7, 37)	r_1	s_1
(67, 13, 36)	$h_1^2 q_1$	$h_0^4 X_3$
(68, 10, 38)	C''	$d_1 t$
(70, 5, 37)	p'	$h_5^2 c_0$
(60, 12, 39)	$h_1^2 X_3$	$h_5 c_0 d_0 e_0$
(70, 14, 37)	τW_1	$\tau P^2 G$

Table 13: Hidden h_2 extensions

(s, f, w)	x	$h_2 \cdot x$
(26, 7, 16)	$h_0 h_2 g$	$h_1^3 h_4 c_0$
(46, 11, 28)	$h_0 h_2 g^2$	$h_1^7 h_5 c_0$
(49, 12, 27)	u'	$\tau h_0 e_0 l$

Table 13: Hidden h_2 extensions

(s, f, w)	x	$h_2 \cdot x$
(50, 11, 28)	$e_0 r$	$Ph_1^3 h_5 c_0$
(52, 12, 29)	v'	$\tau h_0 e_0 m$
(54, 9, 30)	$h_2 B_2$	$h_1 h_5 c_0 d_0$
(57, 16, 31)	Pu'	$\tau h_0 d_0 e_0 j$
(58, 8, 32)	B_6	$\tau e_1 g$
(59, 11, 31)	Q_1	$\tau h_0 B_{21}$
(60, 8, 32)	Q_2	$h_5 k$
(60, 9, 32)	$h_0 Q_2$	$h_0 h_5 k$
(60, 16, 33)	Pv'	$\tau h_0 e_0^2 j$
(62, 13, 34)	$h_0^2 B_{21}$	$Ph_1 h_5 c_0 d_0$
(63, 8, 34)	B_3	$h_5 l$
(63, 15, 38)	$h_0^2 g^3$	$h_1^7 h_5 c_0 e_0$
(63, 16, 35)	$d_0 u'$	$\tau h_0 e_0^2 k$
(64, 7, 34)	$A + A'$	$h_0 A''$
(64, 8, 34)	$h_0(A + A')$	$h_0^2 A''$
(64, 8, 35)	B_7	τd_1^2
(64, 14, 36)	km	$h_1^6 X_1$
(65, 20, 35)	$P^2 u'$	$\tau h_0 d_0^3 i$
(66, 14, 40)	$h_2 g^3$	$h_1^9 D_4$
(66, 16, 37)	$e_0 u'$	$\tau h_0 e_0^2 l$
(67, 15, 36)	U	$h_0 d_0 x'$
(68, 10, 37)	$h_2 c_0$	$\tau d_1 t$
(68, 14, 36)	R_2	$\tau h_1 d_0 x'$
(68, 20, 37)	$P^2 v'$	$\tau h_0 d_0^3 j$
(69, 16, 39)	$e_0 v'$	$\tau h_0 e_0^2 m$
(70, 15, 40)	lm	$h_1^5 c_0 Q_2$

Table 14: Generators for Ext in stems 0–30

element	(m, s, f, w)	description	occurs
h_0	(1, 0, 1, 0)		1
h_1	(1, 1, 1, 1)		1
h_2	(1, 3, 1, 2)		1
h_3	(1, 7, 1, 4)		1
c_0	(5, 8, 3, 5)	$h_1 h_0(1)$	4
Ph_1	(9, 9, 5, 5)		6
Ph_2	(9, 11, 5, 6)		6
d_0	(8, 14, 4, 8)	$h_0(1)^2$	4
h_4	(1, 15, 1, 8)		1
Pc_0	(13, 16, 7, 9)		6
e_0	(8, 17, 4, 10)	$b_{21}h_0(1)$	4
P^2h_1	(17, 17, 9, 9)		12
f_0	(8, 18, 4, 10)	$h_2\nu$	6
c_1	(5, 19, 3, 11)	$h_2h_1(1)$	4
P^2h_2	(17, 19, 9, 10)		12
τg	(8, 20, 4, 11)	τb_{21}^2	6
Pd_0	(16, 22, 8, 12)		6
h_2g	(9, 23, 5, 14)		6
i	(15, 23, 7, 12)	$P\nu$	6
P^2c_0	(21, 24, 11, 13)		12
Pe_0	(15, 25, 7, 14)		6
P^3h_1	(25, 25, 13, 13)		12
j	(15, 26, 7, 14)	$h_0b_{30}h_0(1)^2$	4
h_3g	(9, 27, 5, 16)		6
P^3h_2	(25, 27, 13, 14)		12
k	(15, 29, 7, 16)	$d_0\nu$	6
r	(14, 30, 6, 16)	Δh_2^2	6
P^2d_0	(24, 30, 12, 16)		12

Table 14: Generators for Ext in stems 31–41

element	(m, s, f, w)	description	occurs
h_5	(1, 31, 1, 16)		1
n	(11, 31, 5, 17)	$h_2b_{30}h_1(1)$	4
d_1	(8, 32, 4, 18)	$h_1(1)^2$	4
q	(14, 32, 6, 17)	Δh_1h_3	12
l	(15, 32, 7, 18)	$e_0\nu$	6
P^3c_0	(29, 32, 15, 17)		12
p	(8, 33, 4, 18)	$h_0\nu_1$	6
P^2e_0	(24, 33, 12, 18)		12
P^4h_1	(33, 33, 17, 17)		32
Pj	(23, 34, 11, 18)		6
m	(15, 35, 7, 20)	$g\nu$	6
P^4h_2	(33, 35, 17, 18)		32
t	(12, 36, 6, 20)	$\tau b_{21}^2h_1(1) + h_1^2b_{22}b_{30}$	4
x	(11, 37, 5, 20)	$h_2b_{22}b_{30} + h_2^3b_{40}$	4
e_0g	(16, 37, 8, 22)		6
e_1	(8, 38, 4, 21)	$b_{22}h_1(1)$	4
y	(14, 38, 6, 20)	Δh_3^2	8
P^3d_0	(32, 38, 16, 20)		12
c_1g	(13, 39, 7, 23)		6
u	(21, 39, 9, 21)	Δh_1d_0	6
P^2i	(31, 39, 15, 20)		12
f_1	(8, 40, 4, 22)	$h_3\nu_1$	6
τg^2	(16, 40, 8, 23)		12
P^4c_0	(37, 40, 19, 21)		32
c_2	(5, 41, 3, 22)	$h_3h_2(1)$	4
z	(22, 41, 10, 22)	$\Delta h_0^2e_0$	6
P^3e_0	(32, 41, 16, 22)		12
P^5h_1	(41, 41, 21, 21)		32

Table 14: Generators for Ext in stems 42–53

element	(m, s, f, w)	description	occurs
v	(21, 42, 9, 23)	$\Delta h_1 e_0$	6
$P^2 j$	(31, 42, 15, 22)		12
$h_2 g^2$	(17, 43, 9, 26)		12
$P^5 h_2$	(41, 43, 21, 22)		32
g_2	(8, 44, 4, 24)	b_{22}^2	4
τw	(21, 45, 9, 24)	$\tau \Delta h_1 g$	12
N	(18, 46, 8, 25)	$\Delta h_2 c_1$	6
B_1	(17, 46, 7, 25)	$Y h_1$	8
$h_3 g^2$	(17, 47, 9, 28)		12
u'	(25, 46, 11, 25)	$\Delta c_0 d_0$	6
$P^4 d_0$	(40, 46, 20, 24)		32
Q'	(29, 47, 13, 24)	$P \Delta h_0^2 \nu$	6
$P u$	(29, 47, 13, 25)	$P \Delta h_1 d_0$	6
B_2	(17, 48, 7, 26)	$Y h_2$	8
$P^5 c_0$	(45, 48, 23, 25)		32
v'	(25, 49, 11, 27)	$\Delta c_0 e_0$	6
$P^4 e_0$	(40, 49, 20, 26)		32
$P^6 h_1$	(49, 49, 25, 25)		32
C	(14, 50, 6, 27)	$h_2 x_{47}$	6
gr	(22, 50, 10, 28)	$\Delta h_2^2 g$	6
$P v$	(29, 50, 13, 27)	$P \Delta h_1 e_0$	6
$P^3 j$	(39, 50, 19, 26)		6
G_3	(21, 51, 9, 28)	$\Delta h_3 g$	6
gn	(19, 51, 9, 29)		6
$P^6 h_2$	(49, 51, 25, 26)		32
D_1	(11, 52, 5, 28)	$h_2 x_{49}$	6
$d_1 g$	(16, 52, 8, 30)		6
i_1	(15, 53, 7, 30)	$g \nu_1$	6

Table 14: Generators for Ext in stems 53–60

element	(m, s, f, w)	description	occurs
B_8	(21, 53, 9, 29)	$Y c_0$	8
x'	(24, 53, 10, 28)	$P Y$	6
τG	(14, 54, 6, 29)	$\tau \Delta_1 h_1^2$	6
R_1	(26, 54, 10, 28)	$\Delta^2 h_2^2$	12
$P u'$	(33, 54, 15, 29)	$P \Delta c_0 d_0$	6
$P^5 d_0$	(48, 54, 24, 28)		32
B_6	(17, 55, 7, 30)	$B h_1 h_1(1)$	4
gm	(23, 55, 11, 32)	$g^2 \nu$	6
$P^2 u$	(37, 55, 17, 29)		12
$P^4 i$	(47, 55, 23, 28)		32
gt	(20, 56, 10, 32)		6
Q_1	(26, 56, 10, 29)	$\Delta^2 h_1 h_3$	12
$P^6 c_0$	(53, 56, 27, 29)		32
D_4	(14, 57, 6, 31)	$h_1 b_{21} h_0(1, 2)$	4
Q_2	(19, 57, 7, 30)	$\Delta \nu_1$	6
D_{11}	(21, 57, 9, 31)	$\Delta h_1 d_1$	6
$P v'$	(33, 57, 15, 31)	$P \Delta c_0 e_0$	6
$P^5 e_0$	(48, 57, 24, 30)		32
$P^7 h_1$	(57, 57, 29, 29)		32
D_2	(16, 58, 6, 30)	$h_0 b_{30} h_0(1, 2)$	4
$e_1 g$	(16, 58, 8, 33)		6
$P^2 v$	(37, 58, 17, 31)		12
$P^4 j$	(47, 58, 23, 30)		32
j_1	(15, 59, 7, 33)	$h_1 b_{21} b_{22} b_{31}$	4
B_{21}	(24, 59, 10, 32)	$Y d_0$	8
$c_1 g^2$	(21, 59, 11, 35)		12
$P^7 h_2$	(57, 59, 29, 30)		32
B_3	(17, 60, 7, 32)	$Y h_4$	8

Table 14: Generators for Ext in stems 60–65

element	(m, s, f, w)	description	occurs
B_4	(23, 60, 9, 32)	$Y\nu$	6
τg^3	(24, 60, 12, 35)		12
$h_0 g^3$	(25, 60, 13, 36)		12
D_3	(10, 61, 4, 32)	$h_4 h_0(1, 2)$	4
A	(16, 61, 6, 32)	$h_2 b_{30} h_0(1, 2)$	4
A'	(16, 61, 6, 32)	$h_2 b_{30} h_0(1, 2) + h_0 h_3 b_{31} b_{40}$	4
B_7	(17, 61, 7, 33)	$B h_1 b_{22}$	4
X_1	(23, 61, 9, 32)	Δx	6
H_1	(13, 62, 5, 33)	$h_1(1) h_0(1, 2)$	4
C_0	(20, 62, 8, 33)	$h_2 x_{59}$	6
E_1	(20, 62, 8, 33)	Δe_1	6
B_{22}	(24, 62, 10, 34)	$Y e_0$	8
R	(26, 62, 10, 32)	$\Delta^2 h_3^2$	12
$P^2 u'$	(41, 62, 19, 33)		12
$P^6 d_0$	(56, 62, 28, 32)		32
h_6	(1, 63, 1, 32)		1
C'	(17, 63, 7, 34)	$h_2 b_{40} h_1(1)^2$	4
X_2	(17, 63, 7, 34)	$\tau h_1 b_{21} b_{31}^2 + h_1^2 h_3 b_{31} b_{40}$	4
$h_2 g^3$	(25, 63, 13, 38)		12
$P^3 u$	(45, 63, 21, 33)		12
A''	(14, 64, 6, 34)	$h_0 b_{31} h_0(1, 3)$	4
q_1	(26, 64, 10, 33)	$\Delta^2 h_1 h_4$	32
U	(34, 64, 14, 34)	$\Delta^2 h_1^2 d_0$	12
$P^7 c_0$	(61, 64, 31, 33)		32
k_1	(15, 65, 7, 36)	$d_1 \nu_1$	6
τB_{23}	(24, 65, 10, 35)		8
R_2	(33, 65, 13, 34)	$\Delta^2 h_0 e_0$	12

Table 14: Generators for Ext in stems 65–69

element	(m, s, f, w)	description	occurs
$\tau g w$	(29, 65, 13, 36)		12
$P^2 v'$	(41, 65, 19, 35)		12
$P^6 e_0$	(56, 65, 28, 34)		32
$P^8 h_1$	(65, 65, 33, 33)		34
r_1	(14, 66, 6, 36)	$\Delta_1 h_3^2$	6
τG_0	(17, 66, 7, 35)	$\tau b_{21}^2 h_0(1, 2) +$ $+ h_1 h_3 b_{30} h_0(1, 2)$	4
τB_5	(24, 66, 10, 35)	$\tau b_{21}^2 b_{30}^2 b_{22} +$ $+ \tau h_2^2 b_{21}^2 b_{30} b_{40} + h_1 h_3 b_{30}^3 b_{22}$	4
D_2'	(24, 66, 10, 34)	$P D_2 + \Delta h_0 h_3 x_{35}$	6
$P^3 v$	(45, 66, 21, 35)		12
$P^5 j$	(55, 66, 27, 34)		32
n_1	(11, 67, 5, 36)	$h_3 b_{31} h_2(1)$	4
τQ_3	(13, 67, 5, 35)		12
$h_0 Q_3$	(14, 67, 6, 36)		12
C''	(21, 67, 9, 37)	$g x_{47}$	6
X_3	(21, 67, 9, 36)	$\Delta_1 h_0^2 \nu + \tau \Delta_1 h_1 d_0$	6
C_{11}	(29, 67, 11, 35)	$\Delta^2 c_1$	12
$h_3 g^3$	(25, 67, 13, 40)		12
$P^8 h_2$	(65, 67, 33, 34)		34
d_2	(8, 68, 4, 36)	$h_2(1)^2$	4
G_{21}	(20, 68, 8, 36)	Δg_2	6
$h_2 B_{23}$	(25, 68, 11, 38)		8
G_{11}	(33, 68, 13, 36)	$\Delta^2 h_2 e_0$	12
p'	(8, 69, 4, 36)	$h_0 \nu_2$	6
D_3'	(18, 69, 8, 37)	$h_1 b_{20} b_{31} h_0(1, 3) +$ $+ h_1^3 b_{40} h_0(1, 3)$	4
$h_2 G_0$	(18, 69, 8, 38)		6


Table 14: Generators for Ext in stems 69–70


element	(m, s, f, w)	description	occurs
$P(A + A')$	(24, 69, 10, 36)		6
$h_2 B_5$	(25, 69, 11, 38)		8
τW_1	(33, 69, 13, 36)	$\tau \Delta^2 h_1 g$	12
$P^2 x'$	(40, 69, 18, 36)		12
p_1	(8, 70, 4, 37)	$h_1 \nu_2$	6
$h_2 Q_3$	(14, 70, 6, 38)		12
R'_1	(41, 70, 17, 36)	$\Delta^2 P h_0 d_0$	12
$P^3 u'$	(49, 70, 23, 37)		12
$P^7 d_0$	(64, 70, 32, 36)		32


7. Ext CHARTS


The charts show the cohomology of the motivic Steenrod algebra over $\text{Spec } \mathbb{C}$. Here is a key for reading the charts.

- (1) An element of degree (s, f, w) appears at coordinates (s, f) .
- (2) Black dots indicate copies of $\mathbb{M}_2$.
- (3) Red dots indicate copies of $\mathbb{M}_2/\tau$.
- (4) Blue dots indicate copies of $\mathbb{M}_2/\tau^2$.
- (5) Green dots indicate copies of $\mathbb{M}_2/\tau^3$.
- (6) Vertical lines indicate h_0 multiplications. These lines might be black, red, blue, or green, depending on the τ torsion of the target.
- (7) Lines of slope 1 indicate h_1 multiplications. These lines might be black, red, blue, or green, depending on the τ torsion of the target.
- (8) Lines of slope $1/3$ indicate h_2 multiplications. These lines might be black, red, blue, or green, depending on the τ torsion of the target.
- (9) Red arrows indicate an infinite tower of h_1 multiplications, all of which are annihilated by τ .
- (10) Magenta lines indicate that an extension hits τ times a generator. For example, $h_0 \cdot h_0 h_2 = \tau h_1^3$ in the geometric 3-stem.
- (11) Orange lines indicate that an extension hits τ^2 times a generator. For example, $h_0 \cdot h_0^3 x = \tau^2 h_0 e_0 g$ in the geometric 37-stem.
- (12) Dotted lines indicate that the extension is hidden in the May spectral sequence.
- (13) Squares indicate that there is a τ extension that is hidden in the May spectral sequence. For example, $\tau \cdot P c_0 d_0 = h_0^5 r$ in the geometric 30-stem.









REFERENCES

- [Bl] S. Bloch, *Algebraic cycles and higher K-theory*, Adv. Math. **61** (1986) 267-304.
- [Br1] R. R. Bruner, *The cohomology of the mod 2 Steenrod algebra: A computer calculation*, Wayne State University Research Report **37** (1997).
- [Br2] R. R. Bruner, *Squaring operations in $\text{Ext}_A(F_2, F_2)$* , preprint.
- [DHI] D. Dugger, M. A. Hill, and D. C. Isaksen, *The cohomology of the motivic Steenrod algebra over $\mathbb{R}$* , in preparation.
- [DI] D. Dugger and D. C. Isaksen, *The motivic Adams spectral sequence*, Geom. Topol. **14** (2010) 967-1014.
- [I1] D. C. Isaksen, *The cohomology of motivic $A(2)$* , Homology, Homotopy Appl. **11** (2009) 251-274.
- [I2] D. C. Isaksen, *Some differentials in the motivic Adams spectral sequence over $\text{Spec } \mathbb{C}$* , in preparation.
- [MT] M. Mahowald and M. Tangora, *An infinite subalgebra of $\text{Ext}_A(Z_2, Z_2)$* , Trans. Amer. Math. Soc. **132** (1968) 263-274.
- [Ma1] J. P. May, *The cohomology of restricted Lie algebras and of Hopf algebras; application to the Steenrod algebra*, Ph.D. dissertation, Princeton, 1964.
- [Ma2] J. P. May, *A general algebraic approach to Steenrod operations*, in The Steenrod algebra and its applications (Columbus, 1970), 153-231, Lecture Notes in Mathematics **168**, Springer, 1970.
- [Mi] J. R. Milgram, *Steenrod squares and higher Massey products*, Bol. Soc. Mat. Mexicana **13** (1968) 32-57.
- [T] M. C. Tangora, *On the cohomology of the Steenrod algebra*, Math. Z. **116** (1970) 18-64.
- [V1] V. Voevodsky, *Motivic cohomology with $\mathbb{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. **98** (2003) 59-104.
- [V2] V. Voevodsky, *Reduced power operations in motivic cohomology*, Publ. Math. Inst. Hautes Études Sci. **98** (2003) 1-57.
- [V3] V. Voevodsky, *Motivic Eilenberg-Mac Lane spaces*, Publ. Math. Inst. Hautes Études Sci. **112** (2010) 1-99.